

Antioxidant activities and phenolic contents of bark and leave extracts from Tunisian native tree: *Fraxinus angustifolia* Vahl. subsp. *Angustifolia*

I. TOUHAMI^{1*}, H. GHAZGHAZI¹, H. SELIMI², A. KHALDI¹, H. MAHMOUDI³

¹ University of Carthage. The National Research Institute of Rural Engineering, Water and Forestry. INRGREF. Laboratory of Management and Valorization of Forest Resources. BP 10 Ariana 2080. Tunisia.

² University of Jendouba. Higher Institute of Biotechnology of Beja, ISBB. BP 382-Beja 9000. Tunisia.

³ Université of Tunis El Manar. Unité de Physiologie et Biochimie de la Réponse des Plantes aux Contraintes Abiotiques, Département des Sciences Biologiques, FST, Campus Universitaire, 2092 Tunis, Tunisia.

*Corresponding author: issam_touhami@yahoo.fr

Abstract – The secondary metabolite composition and antioxidant activities of barks and leaves of *Fraxinus angustifolia* Vahl. subsp. *angustifolia* (Oleaceae), Tunisian native tree, from two provenances (Béja and Nefza) were investigated using two solvents extracts (ethanol and distilled water). The highest amounts of polyphenols ($24,84 \pm 0,57$ mg GAE/g DW), flavonoids ($2,71 \pm 0,15$ mg CE/g DW), total tannins ($138,07 \pm 5,77$ mg CE/g DW) and condensed tannins ($68,43 \pm 5,76$ mg CE/g DW) were shown for Béja bark ethanolic extracts. The efficiency of the solvents used to extract phenols from the two organs varied considerably. The antioxidant activity was evaluated by scavenging the radicals 1,1-diphenyl-2-picrylhydrazyl radical (DPPH). The level of antioxidant activity estimated by DPPH test systems was high for Nefza bark ($IC_{50} = 7,12 \pm 0,07$ μ g/ml) and Nefza ethanolic extracts leaves ($IC_{50} = 8,81 \pm 0,20$ μ g/ml). A significant correlation between radical-scavenging capacities of extracts with total phenolic compound was observed. This study was the first report of the antioxidant activity of extracts from leave and bark extracts of *F. angustifolia* in Tunisia. These promising results open the way for further investigations to purify and identify active molecules.

Keywords: *Fraxinus angustifolia*; Oleaceae; leave and bark extracts; bioactive compounds; DPPH method

1. Introduction

Among the various medicinal and culinary tree, some native species are of particular interest because they may be used for the production of raw materials or preparations containing molecules with significant antioxidant capacities and health benefits. Among these species, we listed *Fraxinus angustifolia* Vahl. subsp. *angustifolia* (Oleaceae) as Tunisian native tree. The search for natural substances that can be used as antioxidants in the food, cosmetic and health industries has been intensified (Repetto and Llesuy 2002). In plants, these compounds are mainly represented by polyphenols, flavonoids, and tannins (Lizcano et al. 2010). Polyphenols are a heterogeneous group of secondary metabolites that have a basic structure containing a functional hydroxyl group attached to an aromatic ring. Flavonoids such as anthocyanins, flavonols, and flavones accumulate in the epidermis and protect the plant body from the harmful effects of exposure to ultraviolet light and even herbivores (Manach et al. 2005). In recent years, research has focused on medicinal plants to extract natural and low-cost antioxidants that can replace synthetic additives. Several studies have revealed that all natural antioxidants, especially polyphenols, are able to suppress chronic inflammation and prevent or delay various severe diseases, including cancer, thus playing an important role in the prevention and protection of human tissues from oxidative stress (Mena et al. 2014).

Leaves and samaras were reported to be used in decoctions and infusions as anti-rheumatism and bark against hemorrhoids and fever (Baba-Aissa 1999). In North Algeria, *Fraxinus angustifolia* is used to treat inflammatory diseases like arthritis, rheumatism and gout, which are known to be mediated or promoted by oxidative stress (Ayouni et al. 2016). The bark of *Fraxinus angustifolia* is best known for its use as anti-inflammatory; however, it is also used as antioxidant, diuretic, digestive

and astringent (Atmani et al. 2009). In Morocco, *Fraxinus angustifolia* Vahl. is used for many other traditional purposes to treat pathologies of the digestive system, in dermocosmetology and problems of the nervous system (Fakchich and Elachouri 2014). Therefore, this present study was designed to investigate total phenolic contents, antioxidant activity, free radical scavenging potentials of leaves and bark extracts of *Fraxinus angustifolia* Vahl. subsp. *angustifolia* from two provenances belonging to different bioclimatic areas in Tunisia. The detected variability will be interpreted.

2. Material and Methods

2.1. Plant material

Fraxinus angustifolia Vahl. subsp. *angustifolia* leaf and bark samples were harvested on July 2016 from two provenances: Béja (Latitude 36°50'N; Longitude 9°12'E; mean annual rainfall < 660 mm, sub humid bioclimatic stage) and Nefza (Latitude 37°2'N; Longitude 9°5'E; mean annual rainfall < 725 mm, lower-humid bioclimatic stage). From the region of Béja the main soil characteristics are summarized in a texture of 39,1%, 38,3% and 22,6% of sand, silt and clay respectively, 7,7 of pH and 1,03(S/m) of electrical conductivity. From the region of Nefza the soil texture are 41,3%, 37,7% and 21% of sand, silt and clay respectively, 6.4 of pH and 0,83(S/m) of electrical conductivity. Plant material was identified by Professor Zeineb GHRABI-GAMMAR according to a listed voucher specimen in the herbarium of Department of Botany, INAT (National Agronomic Institute of Tunisia). Samples were air dried and then ground to powder by an electric mill.




Figure 1. Plot distribution map for *Fraxinus angustifolia* Vahl. subsp. *angustifolia* in Tunisia (two provenances from Béja and Nefza)

2.2. Ethanolic extract preparation

For each sample, 1 g of the air-dried aerial parts, finely ground with a blade-carbide grinding (IKAWERKType: A:10), was extracted with 10 ml of pure methanol for 24 h in a water bath shaker maintained at room temperature. The extract was filtered using a Whatman No. 4 filter paper and stored in a brown bottle at 4°C prior to further.

2.3. Aqueous extracts preparation

Water extract: 10 g of the weighed plant leaves powder was soaked in 100 ml of boiled hot water. That mixture was boiled for thirty minutes into a conical flask and put for 24hrs. The extract was filtered using filter paper.

2.4. Colorimetric quantification of phenolics extraction

2.4.1. Determination of total polyphenol content

Phenolic content was assayed using the Folin-Ciocalteu reagent using the method of Lister and Wilson (2001). This method was employed to evaluate the phenolic content of the samples. A calibration curve of gallic acid (ranging from 0.005 to 0.05 mg/ml) was prepared, and the results, determined by the

regression equation of the calibration curve ($y = 10,19x - 0,01$; $R = 0,97$), were expressed as mg gallic acid equivalents per gram dry weight of raw material (mg GAE g^{-1} DW). In this methods, 100 μ l of sample (diluted to obtain absorbance in the range of the prepared calibration curve) were dissolved in 500 μ l (1/10 dilution) of the Folin-Ciocalteu reagent and 1 ml of distilled water. The absorbance of all samples was measured at 760 nm using a HACH UV-Vis spectrophotometer (Hach DR 6000, Germany). Triplicate measurements were taken for each sample.

2.4.2. Determination of total flavonoid content

The amount of flavonoid content was measured using the method described by Dewanto et al. (2002). Total flavonoids were expressed as mg quercetin equivalent per gram DW (mg CE g^{-1} DW), through the calibration curve of quercetin ($y = 46,78x - 0,022$; $R = 0,98$). The calibration curve range was 0-400 μ g ml^{-1} . All samples were analyzed in triplicate.

2.4.3. Determination of condensed and total tannins

Contents of tannins were carried out according to Sun et al. (1998). The concentration of tannins was expressed as mg (+)-equivalent catechin/g DW (mg CE g^{-1} DW). The calibration curve (condensed tannins: $y = 42,15x + 1,620$; $R = 0,99$ and total tannins: $y = 0,0602x - 0,0047$; $R = 0,99$) of catechin was established between 0-250 μ g ml^{-1} . All samples were analysed in triplicate.

2.5. Antioxidant activities

2.5.1. Total Antioxidant Activity (TAA) Assay

The quantification of total antioxidant activity was determined by a colorimetric assay using a method described by Prieto et al. (1999). An aliquot (0.1 ml) of leaves fraction was combined to 1 ml of reagent solution (0.6 M sulfuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate). The tubes were incubated at 95 °C for 90 min. After the mixture had cooled to room temperature; the absorbance of each solution was measured at 765 nm (Anthelie Advanced 2, SECOMAM®) against a blank. The antioxidant capacity was expressed as mg gallic acid equivalent per gram of dry weight (mg GAE/g DW) through the calibration curve of gallic acid ($y = 0,176x - 0,041$; $R = 0,99$).

2.5.2. DPPH Radical-Scavenging Activity

In this part of our work, the in vitro antioxidant activities of leaf and bark extracts have been evaluated based on DPPH test. The DPPH radical scavenging capacity was measured according to Hanato et al. (1988) with modifications (Ghazghazi et al. 2014). The absorbance was measured at 517 nm and corresponded to the ability of extract to reduce the stable radical DPPH to the yellow-colored 2,2 diphenyl-1-picrylhydrazine. The antiradical activity was expressed as IC₅₀ (μ g/ml), the extract dose required to induce a 50 % inhibition. A lower IC₅₀ value corresponds to a higher antioxidant activity of plant extract. The ability to scavenge the DPPH radical was calculated using the following equation:

$$DPPH. \text{ scavenging effect} = [(A_0 - A_1)/A_0] \times 100 \quad \text{Eq (1)}$$

Where A₀ is the absorbance of the control at 30 min, and A₁ is the absorbance of the sample at 30 min. Each phase was analysed in triplicate

2.6. Statistical analysis

Statistical analysis of all data was performed using (IBM® SPSS® Statistics 20). A one-way analysis of variance (ANOVA) was carried out. All data was expressed as mean \pm Standard Error (SE) of three replicates. Correlation between phenolic contents and antioxidant activities was ascertained by Pearson Matrix.

3. Results and discussion

3.1. Total phenolic, flavonoids and tannins contents

The total phenolic, flavonoid, and tannins contents varied significantly among the studied parts in the different extracts of the studied species (Table 1). The concentration of phenolic compounds in *Fraxinus angustifolia* Vahl. subsp. *angustifolia* were evaluated among two studied sites. Barks of Béja, determined in ethanol extract, had the highest contents of polyphenols, flavonoids, total tannins and condensed tannins: $24,84 \pm 0,57$ mg GEA/g DW, $2,71 \pm 0,15$ mg CE/g DW, $138,07 \pm 5,77$ mg CE/g

DW, and $68,43 \pm 5,77$ mg CE/g DW, respectively. We can easily conclude that *F. angustifolia* is poor in flavonoids. These values were much lower than those observed for Algerian *Fraxinus angustifolia* Vahl. (Atmani et al. 2009; Ayouni et al. 2016). For example, Atmani et al. (2009) was found a higher level of total polyphenols, flavonoids and tannins contents in the aqueous extracts of *F. angustifolia* barks 215,7 mg CE/g DW; 4,93 mg Eq Quercetin/g DW; and 177,22 mg Tannic Acid/g DW, respectively. However, Ayouni et al. (2016) reported similar values of flavonoids and tannins contents in aqueous crude extracts of Algerian *F. angustifolia* leaves and bark. While these results are not far from ours, discrepancies can be explained by the genetic difference between the species, the type of soil and microclimate.

Our result exhibited significant differences in content of polyphenols, flavonoids, total and condensed tannins according to the solvent extracting. The ethanolic extract was the richest one as compared to the aqueous fractions from both plant parts (see Table 1). A significant difference ($p < 0.05$) between Béja and Nefza provenances was detected specially for barks part according to ethanolic extract. Likewise, the significant variability between the fractions, in the phenolic compounds, may be attributed to the extracting power of the solvent used and its chemical nature, structure, degree of polymerization and the interaction of these compounds with each other (Saada et al. 2014). Then, the differential accumulation of these compounds between organs should be related to their specific tissues and cells (i.e. mesophyll, epiderm, thickness cuticle, chloroplasts, trichomes). Previous reports have attributed the high presence in barks of these compounds and their low content in leaves to the close interaction between organs and to the different processes of biosynthesis and/or degradation, to the transport involved in the distribution of these polyphenols at the plant level and to the phenological organ growth (Fico et al. 2000; Hudaib et al. 2002).

Table 1. Content of total phenolics, flavonoïdes and tannins in bark and leave extracts of two native tree: *Fraxinus angustifolia* Vahl. subsp. *Angustifolia*. Values are given as mean \pm SE (n = 3). LB: Leaves of Béja; LN: Leaves of Nefza; BB: Barks of Béja; and BN: Barks of Nefza

Plant Part	Aqueous extracts (AE)				Ethanolic extracts (EE)			
	Polyphenols (mg GEA/g DW)	Flavonoids (mg CE/g DW)	Total Tannins (mg CE/g DW)	Condensed Tannins (mg CE/g DW)	Polyphenols (mg GEA/g DW)	Flavonoids (mg CE/g DW)	Total Tannins (mg CE/g DW)	Condensed Tannins (mg CE/g DW)
LB	21,52 \pm 0,88	0,95 \pm 0,02	85,32 \pm 23,76	55,10 \pm 3,33	17,55 \pm 1,44	2,27 \pm 0,07	101,41 \pm 3,33	58,43 \pm 5,77
LN	17,55 \pm 1,84	0,83 \pm 0,03	113,07 \pm 2,89	58,10 \pm 1,45	21,86 \pm 1,15	2,18 \pm 0,13	91,41 \pm 3,33	68,43 \pm 5,77
BB	20,53 \pm 0,66	2,42 \pm 0,13	91,41 \pm 3,33	62,43 \pm 3,05	24,84 \pm 0,57	2,71 \pm 0,15	138,07 \pm 5,77	68,43 \pm 5,77
BN	21,86 \pm 1,15	1,92 \pm 0,03	124,74 \pm 2,40	60,10 \pm 4,41	22,85 \pm 0,57	1,96 \pm 0,13	108,07 \pm 5,77	81,77 \pm 8,82


3.2. Total antioxidant activities

Our results demonstrated that the total antioxidant activity (TAA) of *F. angustifolia* was important from leaves and barks among ethanolic extracts with high values from Nefza barks in comparison to all tested samples ($716,29 \pm 17,64$ mg EAG/g DW).

3.3. DPPH radical scavenging activity

Several antioxidant methods have been proposed to evaluate the free radical scavenging capability of plant materials and to explain antioxidant mechanisms and actions. Among these, free synthetic radical scavenging like DPPH test was used for the evaluation of the total antioxidant behavior of extracts. Ethanol extracts of *F. angustifolia* barks exhibited enhanced TAA (in the order of $716,29 \pm 17,64$ mg EAG/g DW) compared with the other barks extracts (aqueous) using a total antioxidant activity assay (Figure 2a). In fact, Ethanol extracts from leaves and barks of tow provenances of *F. angustifolia* showed the lowest IC₅₀ radical scavenging value, indicating high antioxidant activity with IC₅₀ $7,12 \pm 0,07$ μ g/ml from bark of Nefza (Figure 2b). Atmani et al. (2009) reported similar values with aqueous fraction issued from ethyl acetate and chloroform extractions (IC₅₀ = $10,0$ μ g/ml and IC₅₀ = $10,04$ μ g/ml), respectively. Another investigation involving in medicinal plants showed that the DPPH scavenging activity of the methanol extract of *Fraxinus excelsior* is higher than that of hexane and dichloromethane extracts, suggesting that the hydrogen-donating compounds are more likely to be present in polar solvents (Middleton et al. 2005).

Extracting solvent affected significantly the phenolic compounds and antioxidant activities of several extracts. The most efficient solvents extraction was ethanol extracts may be related to the presence of flavonoid-type compounds and other phenolics. So, the high contents of polyphenol compounds in the studied species contribute to their important antiradical and antioxidative activities. According to Zhang et al. (2011), the antioxidant activity is generally attributed to phenolic and flavonoid compounds in plant extracts.



The Pearson Matrix correlation (Table 3) showed different significance level between measured traits. High positive associations were detected among the measured variables or traits. Therefore, positive correlations were detected between TAA and TT; P and CT. High negative correlations were detected between F, CT, TAA and DPPH.

Indeed, a significant negative correlation between the polyphenol contents and IC₅₀ antioxidant activity values were observed indicating that extracts with highest polyphenol contents show lower IC₅₀ values (Sengul et al. 2009; Khlifi et al. 2011). In the present study, a negative significant correlation ($p < 0.01$), estimated by Pearson Matrix correlation, were observed between the total phenol, flavonoids, condensed tannins contents and TAA activity in different organ extracts and the free radical-scavenging values DPPH (Table 2), which is in agreement with previous findings (Atmani et al. 2009; Ayouni et al. 2016)

	Polyphenols (P)	Flavonoids (F)	Total Tannins (TT)	Condensed Tannins (CT)	TAA activity	DPPH activity
Polyphenols (P)	1	0.405*	0.432**	0.594**	0.342	-0.480*
Flavonoides (F)	0.405*	1	0.081	0.453*	0.447*	-0.570**
Total Tannins (TT)	0.432*	0.081	1	0.229	-0.026	0.138
Condensed Tannins (CT)	0.594**	0.453*	0.229	1	0.646**	-0.628**
TAA activity	0.342	0.447*	0.646**	-0.260	1	-0.662**
DPPH activity	-0.480*	-0.570**	-0.628**	0.138	-0.662**	1

*Correlation is significant at the 0.05 level
 ** Correlation is significant at the 0.01 level

4. Conclusion

The present study provides the first investigation on phenolic contents and antioxidant activities of various extracts from leaves and barks of *Fraxinus angustifolia* Vahl. subsp. *Angustifolia*, native species from Tunisia. Our results clearly showed that the barks was the organ which gives the highest level in polyphenol, flavonoid, total and condensed tannins contents which favor them in industrial use for extraction.

Also, we can conclude that barks ethanolic extracts with higher antioxidant capacity ($IC_{50} = 7,12 \pm 0.07$ $\mu\text{g/ml}$ by DPPH assay and $716,29 \pm 17,64$ mg EAG/g DW by TAA) could be considered as good sources of natural antioxidant. All those results emphasize the importance of the chemical composition of this Tunisian native species.

5. References

- Atmani D, Nassima Chaher N, Meriem Berboucha M, et al (2009)** Antioxidant capacity and phenol content of selected Algerian medicinal plants. Food Chem DOI: 10.1016/j.foodchem.2008.05.077.
- Ayouni K, Meriem BR, Hye KK, et al (2016)** Metabolomic tool to identify antioxidant compounds of *Fraxinus angustifolia* leaf and stem bark extracts. Ind Crops Prod ; <http://dx.doi.org/10.1016/j.indcrop.2016.01.00>.
- Baba-Aissa, F (1999)** Encyclopédie des Plantes Utiles. Flore d'Algérie et du Maghreb, Substances Végétales d'Afrique d'Orient et d'Occident. EDAS, Librairie Moderne, Rouiba, p.368.
- Dewanto V, Wu X, Adom KK, Liu RH (2002)** Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. J Agric Food Chem 50:3010-3014.
- Ghazghazi H, Aouadhi C, Hamrouni S, Mnif W (2015)** Antibacterial, Antifungal and antioxidant activities of Tunisian *Olea europea* ssp. Oleaster fruit pulp and its essential fatty acids. Int J Pharm Pharm Sci Vol 7, Issue 1. (2015) 52-55.
- Hanato T, Kagawa H, Yasuhara T, Okuda, T (1988)** Two new flavonoids and other constituents in licorice root: their relative astringency and radical scavenging effect. Chem Pharm Bull 36, 1090-1097.
- Fakchich, J, Elachouri, M (2014)** Ethnobotanical survey of medicinal plants used by people in Oriental Morocco to manage various ailments. J Ethnopharmacol 154, 76-87.
- Fico, G, Bilia, AR, Morelli, I, Tome, F (2000)** Flavonoid distribution in *Pyracantha coccinea* plants at different growth phases. Biochem Syst Ecol 28, 673-678.
- Hudaib M, Speroni E, Di Pietra AM, Cavrini V (2002)** GC/MS evaluation of Thyme (*Thymus vulgaris* L.) oil composition and variations during the vegetative cycle. J Pharm Biomed Anal 19, 691-700.
- Khlifi D, Hamdi M, El Hayouni A, et al (2011)** Global chemical composition and antioxidant and anti-tuberculosis activities of various extracts of *Globularia alypum* L. (Globulariaceae) leaves. Molecules 16, 10592-10603.
- Lister E, Wilson P (2001)** Measurement of total phenolics and ABTS assay for antioxidant activity (personal communication). Lincoln, New Zealand: Crop Research Institute.
- Lizcano LJ, Bakkali F, Begoña Ruiz-Larrea M, Ignacio Ruiz-Sanz J (2010)** Antioxidant activity and polyphenol content of aqueous extracts from Colombian Amazonian plants with medicinal use. Food Chem vol. 119, no. 4, pp. 1566-1570.
- Manach C, Williamson G, Morand C, et al (2005)** Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. Am J of Clin Nutr, vol. 81, no. 1, (suppl.), pp. 230-242.
- Menaa F, Menaa A, Menaa B (2014)** Polyphenols nano-formulations for topical delivery and skin tissue engineering, in: R.R. Watson (Ed.), Polyphenols Hum Heal Dis p. 65.
- Middleton P, Stewart F, Al-Qahtani S, et al (2005)** Antioxidant, antibacterial activities and general toxicity of *Alnus glutiosa*, *Fraxinus excelsior* and *Papaver rhoeas*. Iran J Pharm Res, 2, 81-86.
- Prieto P, Pineda M, Aguilar M (1999)** Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin. E Anal Biochem 269: 337-341.
- Repetto MG, Llesuy SF (2002)** Antioxidant properties of natural compounds used in popular medicine for gastric ulcers. Braz J Med Biol Res vol. 35, no. 5, pp. 523-534.
- Saada M, Falleh H, Jalleli I, Snoussi M, Ksouri R (2014)** Phenolic profile, biological activities and fraction analysis of the medicinal halophyte *Retama raetam*. S. Afr J of Bot 94: 114-121.
- Sengul M, Yildiz H, Gungor, N, et al (2009)**. Total phenolic content, antioxidant and antimicrobial activities of some medicinal plants. Pak J Pharm Sci 22, 102-106.
- Sun B, Richardo-Da Silvia JM, Spranger I (1998)** Spranger, Critical factors of vanillin assay for catechins and proanthocyanidins, J Agric Food Chem 46, 4267-4274.
- Zhang L, Ravipati AS, Koyyalamudi SR, et al (2011)** Antioxidant and anti-inflammatory activities of selected medicinal plants containing phenolic and flavonoid compounds. J Agric Food Chem 59, 12361-12367.