

Biochemical characterization of the oil of durum wheat germ (Tunisian variety)

M. KAMMOUN¹, N. RASSAA², M. MEJRI-GAÏDA³, A. RICHEL⁴

¹National Institute of Agronomy of Tunis, 43 Avenue Charles Nicole, cité El Mahragene- 1082 Tunis.

²Higher School of Agriculture of Kef, 7100 Boulifa, Le Kef.

³Faculty of Sciences of Tunis, Laboratory of Plant ecology, 9100 El Manar 1, Tunis.

⁴University of Liège, Gembloux Agro-Bio -Tech Department of Chemistry and Bio-industries, Passage of deportees, 2 - 5030 Gembloux Belgium.

*Auteur correspondant: kammounmaroua90@gmail.com

Abstract - The wheat germ oil may constitute an alternative to the misuse of this cereal grain resource. His Tunisian exploitation is limited, at present, in the human and animal feed in the grain form, straw or some transformed products. For the purpose of characterization, the wheat germ oil has been extracted by Soxhlet apparatus from a Tunisian variety of durum wheat (Maali) based on the hexane as a solvent. An analysis by gas chromatography (GC) was then performed to identify and quantify the fatty acids of the oil. Then, the oil has submitted to a second chromatography analysis in liquid phase (HPLC) to quantify the tocopherols. The most important fatty acids constituting have been the two polyunsaturated acids linoleic acid C18:2 (OMEGA 6) and the gamma acid- linoleic C18:3 : 58.24 % and 4.19 % respectively; and the acid mono-unsaturated oleic acid C18:1 (24.49 %) but also some saturated acid: palmitic acid C16:0 with (19.50 %), and the stearic acid C18:0 (1.29%). For this variety of Tunisian durum wheat, the wheat germ oil contains, in addition, 80.27 mg / 100g of α - tocopherols (vitamin E). The wheat germ oil extracted from the Tunisian variety 'Maali' has proved rich in vitamin E and omega-6 ; a better valorisation on the nutritional or pharmaceutical plan is to this title recommended.

Key words: wheat germ, tocopherol, omega-6

1. Introduction

Wheat is among the most important cereal crops worldwide (Morancho, 2000) and particularly in Tunisia (Selma *et al.*, 2005). Food consumption per habitant is, actually, around the 66 kg per year (Terones and Burny, 2012). Semolina is the main product of durum wheat's culture. It constitutes, in fact, 80 to 90 % of the grain mainly at the endosperm (Feillet, 2000). It is essentially composed of starch and gluten.

Moreover, essential minerals (Brandolini and Vidal, 2012), fatty acids (Pomeranz, 1988) and principally vitamins (Giménez *et al.*, 2013) are contained, for the most part, in the germ (Panifili *et al.*, 2003). The grain, as product remains thus under-exploited (Piras *et al.*, 2009). Byproducts such as gru, the bran and germ of wheat are intended, mainly for animal feed. While the extraction of fatty acids from wheat germ can produce an oil for multiple uses: better quality of food for its richness in Omega 6 (Panifili *et al.*, 2003 ;Gelmez *et al.*, 2009), cosmetics owing to the quantities and qualities of Vitamin E or pharmaceutical (Panifili *et al.*, 2003) that can be used to formulate nutraceutical with potential applications for reducing oxidative stress or several other purposes. Wheat germ oil presents, therefore, a co- product of more to enhance further this cereal crop. This oil contains, in fact, essential polyunsaturated fatty acids (Shao *et al.*, 2008) which can't be synthesized by mammals, so essential for the organism. In addition, it is rich in vitamin E and more particularly α -tocopherol (Kumar and Krishna., 2013) which is the most biologically active form. It essentially acts as antioxidants (Gelmez *et al.*, 2009). In this work, a durum wheat germ's oil extracted from a Tunisian variety "Maali" will be, then, characterized.

2. Material and Methods

2.1 Vegetable materials:

Tunisian variety of durum wheat (*Triticum durum*) will be the subject of this study. The variety “Maali” is an improved germoplasm and characterized by large, closed, white and amber grains. It’s a Tunisian obtain from National Institute of Agronomic Research of Tunis (INRAT).

2.2. Extraction of germ:

The germ is separated from the rest of the seed manually after humidification for 2 hours in distilled water. Once extracted, the germs were dried for 24 hours at 50°C and then finely ground

2.3. Extraction of fatty acids:

The oil was extracted by Soxhlet method. The amount of plant material in the cellulose cartridge was 1g. The later is then placed in the Soxhlet extractor to which are attached in the upper part a refrigerant and in the lower part a ground-necked flask containing 70 ml of n-hexane. The cycle was repeated several times until complete exhaustion. At the end of the extraction, the extracted was placed in a flask and dried at 40°C using a vacuum rotary evaporator. Finally it was placed for 30 seconds in an oven at 105 °C so that the solvent had completely evaporated (Dunford and Zhang, 2003).

2.4. Determination of fatty acids:

After preparation of methyl esters of fatty acids, separation by GPC (HP 6890 Series HEWLETT PACKARD) various fatty acids in the sample was performed, according to the FAME 5 method. The chromatograph used in the analysis had the following characteristics: Equilibration 1 min; Maximum temperature 325 °C; Oven program 50 °C for 1 min; Oven N° 1 ;30 °C / min to 150 °C for 0 min; Oven N° 2, 4 °C / min to 240 °C for 10 min (Dong-sun, 1998)

2.5. Tocopherol content:

After sample preparation 100 µl of oil evaporated under dry nitrogen were taken in 1 ml of ACN / MeOH 1/1. An analysis was performed by HPLC (HPLC-UV: 295 nm) by the method (FM: C29H5O02; MM: 430; purity: 99.9%) using the Zorbax 300SB-C18; 4.6 * 150 mm, 3.5 µ; Agilent; and two eluting solvent A: H₂O + 0.5% TFA and B: ACN + MeOH, 1/1 by using DL-alpha Tocopherol Sigma Ref 47783 as standard (Demir and Cetin, 1999)

3. Results

The results given in figure 1, show that the proportions of fatty acids in the durum wheat germ oil vary by constituents. Indeed the highest content is linoleic acid (C18: 2) 58.24%, followed by oleic acid (C18: 1) 24.49% and palmitic acid (C 16: 0) 19.50%. By against, low levels are denoted for the γ -linolenic acid (C18: 3), stearic acid (C18: 0) and Gondolérique acid (C 20: 1) in the range of 4.19%, 1.29% and 0.95%, respectively.

We notice that the rate of α - tocophérol is very important, it is about 802 ppm. In comparison with other vegetable oils, it is highest in wheat germ oil, followed by sunflower oil (671 ppm) and Maize oil (222 ppm). The lowest level (96 ppm) is observed in Olive Oil (Table 1).

Table 1: The levels of α -tocopherol (ppm) in Tunisian durum wheat germ oil compared to other the most consumed vegetable oils (Kamal-Eldin and Andersson, 1997)

Vegetable oils	α - tocopherol in ppm (mg/ kg)
Wheat Germ ¹	802,72
Sunflower ²	671
Peanut ²	141
Soybean ²	116
Olive ²	96
Maize ²	222

¹ Result obtain on Tunisian durum wheat germ oil “Maali”

² Results from works of Kamal-Eldin and Andersson, 1997 [10]




Figure 1: (a) Fatty acids composition for Tunisian durum wheat germ oil (C16 :0 Palmitic Acid, C16: 1 palmitoleic acid, C18: 0 Stearic Acid, C18: 1 Oleic acid, C18: 2 Linoleic acid, C18: 3 γ linolenic acid, C22: 1 erucic acid). (b) Chromatogram (GC) of screening for fatty acids in wheat germ oil.

4. Discussion

The results of the percentages of fatty acids obtained from this analysis are slightly higher than those of Dunford and Zhang (2003) who used the same method of extraction by Soxhlet. Including linoleic acid 58.24% and oleic acid 24.29% against 56.8% and 14.6% respectively which recorded by Dunford and Zhang. In comparison with other vegetable oils, wheat germ oil is the richest in linoleic acid ($\omega 6$) 58.24% followed by Soybean oil 53.7%, Maize oil 44.7% and Olive oil that records the lowest level 9%. This oil is in second place in terms of richness in acid. γ -linolenic 4.19% after 8.1% soybean oil (Kamal-eldin and Andersson, 1997).

Wheat germ oil is, likewise, the richest in α -tocopherol 802.72 ppm relative to other vegetable oils of Olive, Soybean, Maize, Peanut and sunflower. The level of vitamin E which extracted from this variety of wheat “Maali” is higher than that obtained by Kumar and Krishna (2013) on an Indian germoplasm which the level does not exceed 508.59 ppm. This content may, however, vary according to the method and time of extraction from 665 to 2131 ppm (Panifili *et al.*, 2003).

5. Conclusion

This work has helped to highlight the qualities of a durum wheat germ oil, by-product of Tunisian variety “Maali”. The results were very conclusive and showed the richness of the seed on one hand and specifically in fatty acid Omega 6 which is an essential polyunsaturated acid to the body particularly to the nervous system and brain functioning, the prevention of cardiovascular diseases and the formation and maintenance of cell walls and certain immune cells. On the other hand, Tunisian wheat germ oil is revealed rich in α -tocopherol which is the most active form of vitamin E and whose exploitation is widespread in the pharmacological and cosmetics for multiple uses.

6. References

- Brandolini, A., Vidal, A.M.H., (2012).** Wheat germ: not only a by-product. *International Journal of Food Science and Nutrition* 63 (S 1) :71–74.
- Demir C. and Cetin M. (1999)** Determination of tocopherols, fatty acids and oxidative stability of pecan, walnut and sunflower oils. *Deutsche Lebensmittel-Rundschau*, 95 (7) : 278-282.
- Dong-sun L., BONG-Soo N., Sun-Young B., Kun K., (1998)** Characterization of fatty acids composition in vegetable oils by gas chromatography and chemometrics. *Analytica Chimica Acta*, 358 (2) : 163-175.

- Dunford, N. T., & Zhang, M. (2003).** Pressurized solvent extraction of wheat germ oil. *Food Research International*, 36(9-10), 905–909.
- Feillet P.,2000.** Wheat grain composition and use Quae Edition.
- Gelmez, N., Kincal, N. S., &Yener, M. E. (2009).** Optimization of supercritical carbon dioxide extraction of antioxidants from roasted wheat germ based on yield, total phenolic and tocopherol contents, and antioxidant activities of the extracts. *The Journal of SupercriticalFluids*.
- Kamal-eldin, A., &Andersson, R. (1997).** A Multivariate Study of the Correlation Betw
- Kumar, G. S., & Krishna, A. G. G. (2013).** Studies on the nutraceuticals composition of wheat derived oils wheat bran oil and wheat germ oil.
- Morancho, J., (2000).** Production and marketing of durum wheat in the Mediterranean world Option 27: 29-33.
- Panfili, G., Cinquanta, L., Fratianni, A., &Cubadda, R. (2003).** Extraction of Wheat Germ Oil by Supercritical CO 2 : Oil and Defatted Cake Characterization.
- Piras, A., Rosa, A., Falconieri, D., Porcedda, S., Dessì, M. a, &Marongiu, B. (2009).** Extraction of oil from wheat germ by supercritical CO2. *Molecules (Basel, Switzerland)*, 14(7), 2573–81.
- Pomeranz, Y., (1988).** Chemical composition of kernel structures. Wheat: chemistry and technology. **Giménez, I., Herrera, M., Escobar, J., &Ferruz, E. (2013).** Distribution of deoxynivalenol and zearalenone in milled germ during wheat milling and analysis of toxin levels in wheat germ and wheat germ oil. *Food Control*.
- Selma, A., Ben-Salem, M., ben Naceur, M etzid, E.(2005).** The cereal in Tunisia: Production, due to drought and resistance mechanisms. *Drought*. 16 (3): 255-229.
- Shao, P., Sun, P., & Ying, Y. (2008).** Response surface optimization of wheat germ oil yield by supercritical carbon dioxide extraction. *Food and Bioproducts Processing*, 86(3), 227–231.
- Terones Gavira .Fr ; Burny Ph., (2012)** White Paper 'cereals'; Evolution of the world wheat market in the fifty last years. ULg Gembloux Agro Bio Tech and CRA-W Gembloux. Volume I., 97-158.